

EGG YOLK TELLURITE EMULSION 50%

TYPICAL FORMULA (per litre)

Fresh egg yolk470 ml
Saline solution.....470 ml
Potassium Tellurite 3,5%... 70 ml

INTENDED USE

Liquid supplement for the differentiation of *Staphylococcus aureus*. Egg Yolk Tellurite Emulsion should be used for the supplementation of Baird Parker Agar Base. It is used for the detection of lecithinase enzyme activity and for the potassium tellurite reduction test. For the details of the procedure see the technical sheet of Baird Parker Agar Base (cat. N° 401116)

DIRECTIONS

Baird Parker Egg Yolk Tellurite Agar: 50 ml Egg Yolk Tellurite Emulsion + 1000 ml Baird Parker Agar Base (REF 401116) autoclaved and cooled to 45-50°C

STORAGE

Store at 2-8° - When stored as directed the supplement remains stable until the expiry date shown on the label. Do not use beyond stated expiry date. Separation may occur on storage. This occurrence is a normal one and doesn't affect the quality of the product. Mix well before the use

PRECAUTIONS AND WARNING

For *in vitro* diagnostic use only. The supplement should be used only by adequately trained personnel with knowledge of microbiological techniques in the laboratory.
Consult the material safety data sheet before the use.

PACKAGING

42111602	Egg Yolk Tellurite Emulsion 50%,	50 ml
42111604	Egg Yolk Tellurite Emulsion 50%,	100 ml
42111603	Egg Yolk Tellurite Emulsion 50%,	200 ml