

OXYTETRACYCLINE ANTIMICROBIC SUPPLEMENT GENTAMICIN ANTIMICROBIC SUPPLEMENT

VIAL CONTENTS (for 500 ml of medium)

Oxytetracycline Antimicrobial Supplement:

Oxytetracycline HCl.....50 mg

Gentamicin Antimicrobial Supplement:

Gentamicin Sulphate.....25 mg

DIRECTIONS

Aseptically reconstitute one vial of Oxytetracycline Antimicrobial Supplement and one vial of Gentamicin Antimicrobial Supplement with 5 ml of sterile distilled water. Prepare 500 ml of OGYE Agar Base, autoclaved and cooled to 50°C. Add the contents of one vial of Oxytetracycline Antimicrobial Supplement and, if required by laboratory procedures, add also the contents of one vial of Gentamicin Antimicrobial Supplements. Mix well and distribute into sterile Petri dishes..

INTENDED USE

Once the above described lyophilised supplements have been added to the base media, they can be used for the isolation of yeasts and moulds. For the details of the procedures see the technical sheets of OGYE Agar Base (401838)

STORAGE

Store at 2-8° - When stored as directed the supplements remain stable until the expiry date shown on the label. Do not use beyond stated expiry date.

PRECAUTIONS AND WARNING

The supplements should be used only by adequately trained personnel with knowledge of microbiological techniques in the laboratory.

Consult the material safety data sheet before the use.

PACKAGING

4240004 Gentamicin Antimicrobial Supplement 10 vials, each for 500 ml of medium

4240000 Oxytetracycline Antimicrobial Supplement 10 vials, each for 500 ml of medium