

## KANAMYCIN-POLYMYXIN B ANTIMICROBIC SUPPLEMENT

**VIAL CONTENTS** (for 500 ml of medium)

Kanamycin.....6 mg  
Polymyxin B Sulphate.....15.000 IU

**DIRECTIONS**

**SFP Agar:** aseptically reconstitute the contents of one vial of Kanamycin-Polymyxin B Antimicrobial Supplement with 5 ml of sterile distilled water. Prepare 500 ml of Clostridium Perfringens Agar Base, cooled to 50°C. Add the supplement, mix well and distribute into sterile Petri dishes.

The medium may be prepared also by adding 50 ml of Egg Yolk Emulsion 50% (code 42111601) to 450 ml of selective medium, cooled to 50°C.

**INTENDED USE**

For the selective isolation and enumeration of *Clostridium perfringens* in foods. For the details of the procedures see the technical sheets of Clostridium Perfringens Agar Base, cat. N° 401307.

**STORAGE**

Store at 2-8° - When stored as directed the supplement remains stable until the expiry date shown on the label. Do not use beyond stated expiry date.

**PRECAUTIONS AND WARNING**

For *in vitro* diagnostic use only. The supplement should be used only by adequately trained personnel with knowledge of microbiological techniques in the laboratory.

Consult the material safety data sheet before the use.

**PACKAGING**

**4240005 Kanamycin-Polymyxin B Antim. Supplement, 10 vials, each for 500 ml of medium**