

BLASER WANG ANTIMICROBIC SUPPLEMENT

VIAL CONTENTS (for 500 ml of medium)

Amphotericin B	1.0 mg
Cefalotin	7.5 mg
Trimethoprim	2.5 mg
Vancomycin	5.0 mg
Polymyxin B	1250 IU

DIRECTIONS

Aseptically reconstitute the contents of one vial with 5 ml (or 50 ml) of sterile distilled water. Prepare 500 ml (or 5 l) of Columbia Blood Agar Base (cat. N° 401136) or Brucella Medium Base or Blood Agar Base N° 2, autoclaved and cooled to 50°C. Add the supplement together with 50 (or 500) ml of defibrinated sheep blood or 25 (or 250 ml) of lysed horse blood and the contents of one vial of Campylobacter Growth Supplement (cat. N° 4240021). Mix well and pour into sterile Petri dishes.

INTENDED USE

The selective supplement here described is used for the preparation of plating media for the isolation of *Campylobacter* in clinical and other specimens. For the details of the procedures see the Campylobacter Blood Agar Base technical sheet (4012852)

STORAGE

Store at 2-8° - When stored as directed the supplement remains stable until the expiry date shown on the label. Do not use beyond stated expiry date.

PRECAUTIONS AND WARNING

For *in vitro* diagnostic use only. The supplement should be used only by adequately trained personnel with knowledge of microbiological techniques in the laboratory. Consult the material safety data sheet before the use.

PACKAGING

4240015 Blaser Wang Antimicrobial Supplement, 10 vials, each for 500 ml of medium