

LISTERIA PALCAM ANTIMICROBIC SUPPLEMENT

VIAL CONTENTS (for 500 ml of medium)

Polymyxin B	5.0 mg
Ceftazidime	10.0 mg
Acriflavine HCl	2.5 mg

DIRECTIONS

Dissolve the contents of one vial with 5 ml of sterile distilled water under aseptic conditions. Add to 500 ml of Listeria PALCAM Agar Base (cat. N° 401604), autoclaved and cooled to 50°C. Mix well and pour into sterile Petri dishes.

INTENDED USE

The selective supplement here described is used for the preparation of OXFORD plating medium for the isolation of *Listeria* spp. from food, environmental or clinical specimens. For the details of the procedure, see the technical sheet of Listeria PALCAM Agar Base (cat. N° 401604).

STORAGE

Store at 2-8° - When stored as directed the supplement remains stable until the expiry date shown on the label. Do not use beyond stated expiry date.

PRECAUTIONS AND WARNING

For *in vitro* diagnostic use only. The supplement should be used only by adequately trained personnel with knowledge of microbiological techniques in the laboratory.
Consult the material safety data sheet before the use.

PACKAGING

4240042 Listeria PALCAM Antimicrobial Supplement, 10 vials, each for 500 ml of medium